

Volume Conservation, Dark Matter, Dark Energy, and Gravitation

Preprint Revision 0.91

Jeffrey Chamberlain \ `jeffrey.d.chamberlain@klingturfang.net`

October 25, 2026

Abstract

We propose that through a non-metrizable topology $\mathcal{Q}$, there exists a non-classical inseparability between points on the same spacelike hypersurface. We posit that energy concentrations displace spatial volume to induce a volume-conserving potential across $\mathcal{Q}$, from which we derive an equation for the flow of 3-volume along the potential gradient. From this we make a testable prediction of the cosmological distribution of dark energy and show that the gravitational coupling is proportional to the universe's global capacity for absorbing volumetric flow. We close with a derivation of an upper limit for the universe's total energy, and a conjecture that the volumetric pressure upon 3-space asymptotically approaches the Planck volume as the energy density approaches infinity.

1 Introduction

We begin with the conjecture that nature's wave-particle duality[1–4] reflects an intrinsic property of its topological structure, wherein the points of space exist on two topologies simultaneously. This bitopology consists of the familiar metrizable 3-space manifold of our daily experience conjoined with an indiscrete topology on which all points are inseparable.

2 Topological Framework

We take inspiration from the BSSN formalism[5, 6] and foliate the 4-dimensional manifold of spacetime $\mathcal{M}$ into a family of spacelike hypersurfaces $\{\Sigma_i\}$. Let Σ_t be one such hypersurface, $\mathbb{S}$ the set of all points in Σ_t , and γ_{ij} the spatial metric projected on Σ_t from $\mathcal{M}$'s metric tensor, $g_{\mu\nu}$. Decompose γ_{ij} into a dimensionless unimodular component $\bar{\gamma}_{ij}$ and a volume 0-form Ω_t on Σ_t , such that

$$\gamma_{ij} = \Omega_t^{\frac{2}{3}} \bar{\gamma}_{ij} \quad (1)$$

and

$$\gamma = \Omega_t^2. \quad (2)$$

Since our purpose is to ground a theoretical volume conservation principle in differential geometry, we have replaced BSSN's dimensionless conformal factor with a volume form and instead stripped $\bar{\gamma}_{ij}$ of its dimensions.

Define two metric-free topologies over the points of $\mathbb{S}$. The first is a smooth manifold $\mathcal{V}$, with volume 3-form

$$\omega = \Omega_t \omega_0, \quad (3)$$

$\mathcal{V}$ has no metric, and thus lacks $\bar{\gamma}_{ij}$'s measure for distance, angles and area, but is related to $\mathcal{M}$ through

$$\Sigma = (\mathcal{V}, \bar{\gamma}_{ij}). \quad (4)$$

The second is an indiscrete topology $\mathcal{Q}$ on which the only non-empty open set is the set of all points of $\mathbb{S}$. To aid intuition, imagine a relationship between these topologies through a degenerate inner product on $\mathcal{V}$'s tangent bundle that renders points inseparable,

$$\langle u, v \rangle_{\mathcal{Q}} = 0, \forall u, v \in T_p \mathcal{V} \text{ and } p \in \mathcal{V}. \quad (5)$$




Figure 1: Bitopological Framework

Let the full topological structure on the points of space be the bitopology

$$\mathcal{U} \equiv (\mathbb{S}, \mathcal{V}, \mathcal{Q}), \quad (6)$$

or, with the unimodular metric inserted and leaving $\mathbb{S}$ implied,

$$\Upsilon \equiv (\Sigma, \mathcal{Q}). \quad (7)$$

To stitch the spacelike foliation back together we can write $\Upsilon_t = (\Sigma_t, \mathcal{Q}_t)$, and define

$$\mathcal{M} \equiv \{\Sigma_i, \mathcal{Q}_i\}, \quad (8)$$

visualizing $\mathcal{M}$ as a sequence of Υ_t progressing in successive epochs of cosmic time.

2.1 Classical Law

With this adjustment to $\mathcal{M}$'s definition, we posit that non-local quantum entanglement[7–11], tunnelling[12–14], and wave function measurement effects—first framed as a paradox by Einstein, Podolsky, and Rosen[15, 16] yet shown to be inevitable by Bell[17]—avoid violation of relativistic locality by allowing non-classical mechanics through $\mathcal{Q}$'s topological inseparability.

We define non-classical mechanics as those invariant with respect to $\bar{\gamma}_{ij}$. The bitopology $(\mathcal{V}, \mathcal{Q})$ from eq. (6) is the natural space for analyzing such mechanics, where freedom from the constraints of the metric enables effects to operate through $\mathcal{Q}$'s indiscrete connectivity to points anywhere on the same spacelike hypersurface.

Conversely, classical mechanics are those dependent on a geometric measure. Classical effects are subject to the constraints of Σ 's $\bar{\gamma}_{ij}$ in eq. (4). For these effects connectivity through $\mathcal{Q}$ is impassable, as a non-metrizable topology is fundamentally incapable of providing the necessary metric.

We use this definition of a non-classical mechanic to build our volume conservation proposal into an experimentally testable hypothesis. A deeper exploration of the implications to quantum and classical theory is a topic for future work.

2.2 Volume Conservation

We conjecture that $\mathcal{V}$ has constant volume

$$\mathbb{V} \equiv \int_{\mathcal{V}} \omega = \int_{\mathcal{V}} \Omega_t \omega_0, \forall t, \quad (9)$$

and propose that the volume form's conservation mechanism is non-classical and therefore must have no dependence on $\bar{\gamma}_{ij}$. When volumetric pressure requires local deformations in Ω_t to reassign elsewhere on $\mathcal{V}$, the effect operates through the same topology $\mathcal{Q}$ that governs quantum measurement, tunneling, and entanglement.

As unintuitive as a volume conservation mechanism that reassigns units of volume across spacelike intervals may seem, there is related theoretical precedent. A non-metrizable $\mathcal{Q}$ as the topological enabler of such behavior has explanatory power comparable to that of a wormhole[18] but through proposing an exotic topology rather than exotic spacetime curvature. Though before developing our volume conservation mechanic's formal structure, we present a motivational thought experiment.

3 Thought Experiment




Figure 2: Volume Displacement

Start with a vacuum spacetime into which we insert a sphere of energy E with density ρ and radius r at point x^i (fig. 2).

Following the series expansions in [19], our existing relativity theory expects the energy density around x^i to compress the local volume according to

$$\Delta\Omega \propto -\frac{8\pi r^2 G}{15c^2} \rho + \mathcal{O}(r^4). \quad (10)$$

For our thought experiment, with volume conserved, this displaced volume must be offset with a corresponding increase elsewhere in x^i 's spatial slice, as shown in (fig. 2).

3.1 Residual Stress

Imagine some time to pass, remove E , and consider what happens to the stress energy tensor. We would normally observe that the contracted Bianchi identities, $\nabla_\mu G^{\mu\nu} = 0$, require $\nabla_\mu T^{\mu\nu} \rightarrow 0$ to conserve energy locally. But in our thought




Figure 3: Residual Stress

experiment, since this requires that volume around x^i increase to its previous state, we have a contradiction. The volume has already been reassigned globally to preserve $\mathbb{V}$ and thus, the previous volume cannot be so easily restored (fig. 3). What occurs instead is a form of spacetime plasticity reminiscent of that suggested by Feoli[20], where the Bianchi identities require that residual non-zero stress remain amongst the stress-energy terms to offset the unrestored volume.




Figure 4: Dark Matter

Now imagine inserting a unit of energy $E_{1/2}$ at x^i that is equivalent to E except that its energy density is $\frac{\rho}{2}$. Inserting $\rho/2$ into eq. (10) gives the volume pressure from $E_{1/2}$ at x^i ,

$$\Delta\Omega \propto -\frac{1}{2} \cdot \frac{8\pi r^2 G}{15c^2} \rho + \mathcal{O}(r^4). \quad (11)$$

Since this is strictly less than the volume deformation left behind by E from eq. (10), $E_{1/2}$ will settle into E 's residual gravity well with no additional deformation (fig. 4).

3.2 Global Smoothing




Figure 5: Dark Energy

Finally, imagine reinserting E at some y^i such that the separation between x^i and y^i on γ_{ij} is greater than $2r$. A local deformation of volume will form at y^i , and the displaced volume will reassign across y^i 's spatial slice to preserve $\mathbb{V}$.

Most importantly, residual stress around x^i will reduce (fig. 5).

4 Volumetric Potential

Let ζ be a strictly monotonically decreasing, positive-definite smooth function on $\mathbb{R}_{\geq 0}$. Assume energy displaces volume and that for each $q \in \mathcal{V}$, $(\zeta \circ \rho)(q)$ returns the maximum volume allowed at q .

We call ζ the **energy-volume relation**, and define the composition

$$\Omega_\rho \equiv \zeta \circ \rho \quad (12)$$

to be the **volumetric limit**; a positive definite 0-form on $\mathcal{V}$ satisfying $\text{Image}(\Omega_\rho) \subset (0, \zeta(0)]$. Define its maximum,

$$\Omega_{\max} \equiv \zeta(0). \quad (13)$$

to be the **vacuum volume limit**.

4.1 Volumetric Gradient

In a region where Ω_ρ drops below the local volume Ω_t , the volumetric pressure within the region will reduce Ω_t to Ω_ρ by causing $\Omega_t - \Omega_\rho$ units of volume to flow out.

An integral across all such regions yields the **volumetric gradient** for a spacelike foliation Σ_t ,

$$\Delta\Omega = - \int_{\mathcal{V}_-} (\Omega_\rho - \Omega_t) \omega_0, \quad (14)$$

where $\mathcal{V}_- \equiv \{q \in \mathcal{V} \mid \Omega_t(q) > \Omega_\rho(q)\}$.

Define the **volumetric potential** to be

$$\Phi \equiv \Omega_\rho - \Omega_t, \quad (15)$$

and complete the partition of $\mathcal{V}$ with $\mathcal{V}_+ \equiv \{q \in \mathcal{V} \mid \Omega_t(q) < \Omega_\rho(q)\}$ and $\mathcal{V}_0 \equiv \{q \in \mathcal{V} \mid \Omega_t(q) = \Omega_\rho(q)\}$, as shown in fig. 6.




Figure 6: Volumetric Potential

Then we can write

$$\mathcal{V} = \mathcal{V}_- \cup \mathcal{V}_+ \cup \mathcal{V}_0, \quad (16)$$

and summarize our partition of $\mathcal{V}$ succinctly:

- $\mathcal{V}_-$ is the set of all points where $\Phi < 0$. Volume here exceeds the volumetric limit from eq. (12), inducing a volumetric gradient that directs flow away.
- $\mathcal{V}_+$ is the set of all points where $\Phi > 0$. Here, with volume below the volumetric limit, there is excess volumetric potential. The volumetric gradient flows toward these regions with a magnitude proportional to the product of $\Delta\Omega$ with a probability distribution W that varies proportional to the potential Φ . We will formally define W in section 4.2.
- $\mathcal{V}_0$ is the set of all points where $\Phi = 0$. Volume here is at its limit. No volume is flowing away and no volume is flowing in.

We propose that the volumetric flow follows eq. (14)'s gradient from the universe's $\mathcal{V}_-$ partition to its $\mathcal{V}_+$ partition as instantaneously as the Planck time allows, with volume continuously tunneling across spacelike separations through $\mathcal{Q}$. This yields

$$\frac{\Delta\Omega}{\Delta t} = -\frac{1}{t_P} \int_{\mathcal{V}_-} (\Omega_\rho - \Omega_t) \omega_0. \quad (17)$$

Note that this flow does not require anything to be entangled. We only hypothesize that the volumetric potential's gradient is applied across the same non-metrizable topology through which entanglement is maintained, and that this alone is sufficient to enable the tunneling of displaced volume.

For each $q \in \mathcal{V}_-$, the negative outflow is thus Φ / t_P , and for each $q \in \mathcal{V}_+$, the positive inflow is

$$\frac{(W \circ \Phi)}{\int_{\mathcal{V}_+} (W \circ \Phi) \omega_0} \frac{1}{t_P} \int_{\mathcal{V}_-} (-\Phi) \omega_0, \quad (18)$$

where W is a positive definite smooth function on $\mathbb{R}_{>0}$ that weights the distribution of the flow.

4.1.1 Convergence

Before deriving W , we consider the two limits where eq. (18) does not converge. The first is the limit where the volumetric gradient from eq. (14) is unbounded. In this limit, $\mathcal{V}_-$ cannot be compact because $\int_{\mathcal{V}_-} (-\Phi) \omega_0$ is unbounded only if $\int_{\mathcal{V}_-} \omega_0$ is unbounded. Furthermore, we can show that every point of $\mathcal{V}_-$ must have non-zero energy density if we borrow an equation from section 5.

Proof: In eq. (28) we see that if Ω_t ever goes above $\Omega_{\max}$, it will drop back down to $\Omega_{\max}$ within a $\Delta t = 5.39 \cdot 10^{-44}$ seconds (the Planck time, t_P). As a result, since all points in $\mathcal{V}_-$ have the property that $\Omega_t > \Omega_\rho$, we can conclude that Ω_ρ must be strictly less than $\Omega_{\max}$ across $\mathcal{V}_-$. Thus, since ζ 's strict monotonicity

means $\zeta(\rho) = \Omega_{\max}$ only when $\rho = 0$, we conclude that all points in $\mathcal{V}_-$ have $\rho > 0$.

From this it follows that $\mathcal{V}_-$ is non-compact, only if the total energy of the universe $\mathbb{E}$ is unbounded. We defer to Archimedes [21] and propose $\mathbb{E}$ to be finite, and thus conclude that $\mathcal{V}_-$ is compact.

The second is the limit where the integral in the denominator goes to zero. Since $W \circ \Phi$ is positive definite across $\mathcal{V}_+$, we know that $\int_{\mathcal{V}_+} (W \circ \Phi) \omega_0 = 0$ only if $\mathcal{V}_+ = \emptyset$. This represents a universe with no capacity to receive volumetric flow. If this were possible, the volume of 3-space would be infinitely rigid and unable to deform in response to energy. Since it is empirically well established that space does bend in the presence of energy [22, 23], it is therefore safe to assume that the integral over $\mathcal{V}_+$ is non-zero.

Though, as discussed in more detail in section 5.5.1, we will later see that experimental observation of volumetric flow will verify the compactness of both $\mathcal{V}_+$ and $\mathcal{V}_-$.

4.2 Volumetric Flow Distribution

The volumetric potential Φ is a 0-form with $\Phi(\mathcal{V}) \subseteq (-\Omega_{\max}, \Omega_{\max})$. Since, by definition, $\Phi(\mathcal{V}_0) = \{0\}$ and $\Phi(\mathcal{V}_+) \subseteq (0, \Omega_{\max})$, it follows from Φ 's smoothness that $\mathcal{V}_+$'s boundary, $\mathcal{S} = \mathcal{V}_+ \setminus \mathcal{V}_+^\circ$, satisfies $\mathcal{S} \subseteq \mathcal{V}_0$. We can thus assume that for any sequence $\{q_i\} \subset \mathcal{V}_+$ such that $q_i \rightarrow \mathcal{S}$, it is also true that $\Phi(q_i) \rightarrow 0^+$ and $(W \circ \Phi)(q_i) \rightarrow 0^+$.

We expect that the weight distribution, $W \circ \Phi$, from eq. (18) will direct volumetric flow to all of $\mathcal{V}_+$, but should otherwise be repulsed by the boundary $\mathcal{S}$. Take this to mean that W should be infinitely flat as $\Phi \rightarrow 0^+$. The canonical infinitely flat function is Cauchy's example [24] $\{e^{-1/x^2}$ for $x \neq 0$, 0 for $x = 0\}$, but for our purposes infinitely flat is required only for $x \rightarrow 0^+$, and so $e^{-1/x}$ is sufficient.

This suggests W should include a term $e^{-b/\Phi}$, for some constant b with units of volume. Furthermore, since Φ is monotonically decreasing with a volumetric limit Ω_ρ , set by an energy-density ρ that cannot resolve a volume displacement with a granularity finer than the Planck volume ℓ_P^3 , infer $b = \ell_P^3$, and write

$$W \propto e^{-\ell_P^3/\Phi} \quad (19)$$

to cover the behavior of the weighting function at points near the boundary.

An additional term is required to describe W 's behavior in $\mathcal{V}_+$'s interior, where the volumetric flow should be attracted to regions proportionally with how close Φ is to the vacuum volume limit $\Omega_{\max}$. To provide the required attraction, assume any positive exponential term $e^{f(\Phi)}$, such that f increases polynomially with Φ across $\Phi \in (0, \Omega_{\max})$, mathematically suitable, but for our purposes assume a linear f sufficient and propose

$$W \propto e^{a \Phi} \quad (20)$$

to cover the weighting function behavior at points near its maximum. Units of inverse volume are required for the constant a . To ensure consistent scaling regardless of the specific value nature sets for the volumetric limit of the vacuum $\Omega_{\max}$, deduce that $a = 1/\Omega_{\max}$, and write our complete weighting function as

$$W = e^{\Phi/\Omega_{\max}} \cdot e^{-\ell_P^3/\Phi} . \quad (21)$$

We can now write the local volume displacement from eq. (18) as

$$\frac{\Delta\Omega}{\Delta t} = \frac{\exp\left(\frac{\Phi}{\Omega_{\max}} - \frac{\ell_P^3}{\Phi}\right)}{\int_{\mathcal{V}_+} \exp\left(\frac{\Phi}{\Omega_{\max}} - \frac{\ell_P^3}{\Phi}\right) \omega_0} \frac{1}{t_P} \int_{\mathcal{V}_-} (-\Phi) \omega_0 \quad (22)$$

and derive a complete expression for $\partial_t \Omega$.

4.3 Volumetric Flux

For $q \in \mathcal{V}_- \cup \mathcal{V}_0$ where $\Phi \leq 0$, volume flows away at the rate

$$\frac{\partial\Omega}{\partial t} = \frac{1}{t_P} \Phi \quad (23)$$

where the integral of $(-\partial_t \Omega)$ across the non-zero region

$$\frac{1}{t_P} \int_{\mathcal{V}_-} (-\Phi) \omega_0 \quad (24)$$

computes the total **volumetric flux** for the spacelike hypersurface Σ_t .

For $q \in \mathcal{V}_+$, where $\Phi > 0$ indicates excess volumetric capacity, the volume flows in according to

$$\frac{\partial\Omega}{\partial t} = \frac{\exp\left(\frac{\Phi}{\Omega_{\max}} - \frac{\ell_P^3}{\Phi}\right)}{\int_{\mathcal{V}_+} \exp\left(\frac{\Phi}{\Omega_{\max}} - \frac{\ell_P^3}{\Phi}\right) \omega_0} \frac{1}{t_P} \int_{\mathcal{V}_-} (-\Phi) \omega_0 . \quad (25)$$

This describes a distribution for the universe's expansion rate that is testable against cosmological observations. In the most isolated near-vacuum regions of intergalactic space, where variability in the volumetric potential Φ should be small, eq. (25) predicts an expansion rate that is close to constant across points on the same spacelike hypersurface Σ_i —varying smoothly along timelike trajectories orthogonal to $\mathcal{M}$'s spacelike foliation (fig. 1). While across points on

the same Σ_i , the expansion rate varies in proportion with how long it has been since energy densities were high in the region—which means along the rotational and translational past histories of large cosmological mass concentrations.

Finally, notice that our derivation preserves the continuity of space. We assumed a quantized volumetric flux with the $e^{-\ell_P^3/\Phi}$ term in W . But it was the classical demand that integrals not blow up to infinity in their ultraviolet limit[25]—that required energy-densities to be discrete, and thus led to W 's ultraviolet limit. We propose that the manifold $\mathcal{V}$ remains smooth while volume is displaced in quanta proportional to the quantized energetic pressure.

5 Vacuum Spacetime $\mathring{\mathcal{V}}$

In spacetime regions where the energy density ρ approaches vacuum, the volumetric limit Ω_ρ approaches $\Omega_{\max}$, and for vacuum spacetime the volumetric potential from eq. (15) becomes

$$\Phi = \Omega_{\max} - \Omega_t. \quad (26)$$

To organize investigation of $\mathcal{V}$'s global vacuum state, we partition $\mathring{\mathcal{V}}$ into subregions based on the value of Φ , as we did previously with $\mathcal{V}$.




Figure 7: Vacuum Spacetime

5.1 Vacuum State $\mathring{\mathcal{V}}_-$

Definition: $\mathring{\mathcal{V}}_- \equiv \{q \in \mathring{\mathcal{V}} \mid \Omega_{\max} < \Omega_t(q)\}$.

This vacuum region defines an overstrained vacuum. We know from eq. (17) that each point $q \in \mathring{\mathcal{V}}_-$ will experience a volumetric outflux equal to

$$\frac{1}{t_P} (\Omega_{\max} - \Omega_t) \quad (27)$$

that ensures $\Omega_{t+t_P} = \Omega_{\max}$. As a result, for any non-empty $\mathring{\mathcal{V}}_-$ vacuum state,

$$\mathring{\mathcal{V}}_- \rightarrow \emptyset, \text{ as } \Delta t \rightarrow t_P. \quad (28)$$

This shows that no region of 3-space can be in this vacuum state for longer than the Planck time at least so long as $\mathring{\mathcal{V}}_+ \neq \emptyset$, and $\int_{\mathring{\mathcal{V}}_+} (W \circ \Phi) \omega_0$ is sufficiently large to accept the outflux.

5.1.1 When $\mathring{\mathcal{V}} \rightarrow \mathring{\mathcal{V}}_-$

For a vacuum spacetime with $\mathring{\mathcal{V}} = \mathring{\mathcal{V}}_-$ (the top line in fig. 8), we would have $\mathring{\mathcal{V}}_+ = \emptyset$ and a contradiction. There would be volumetric outflux at every point, but no region able to accept the flow. This is an example of the previously discussed limit case of eq. (18) where the denominator goes to zero. It represents a spatial fabric stretched so tight it should tear. We presume a spacetime with $\mathring{\mathcal{V}} = \mathring{\mathcal{V}}_-$ non-physical, or, at best, if physical, representative of a pre-gravitational phase of the volume form Ω_t involving physics beyond the scope of this paper.

5.2 Vacuum State $\mathring{\mathcal{V}}_0$

Definition: $\mathring{\mathcal{V}}_0 \equiv \{q \in \mathring{\mathcal{V}} \mid \Omega_{\max} = \Omega_t(q)\}$.

Across these vacuum regions, $\Phi = 0$. There is no volumetric outflux and no capacity for accepting volumetric influx.

5.2.1 When $\mathring{\mathcal{V}} \rightarrow \mathring{\mathcal{V}}_0$

A vacuum spacetime with $\mathring{\mathcal{V}} = \mathring{\mathcal{V}}_0$ is stable, but static, with a constant volume of $\Omega_{\max}$ at every point (the middle line in fig. 8). With no sink to receive volumetric flow, any energy inserted would be unable to displace volume, implying that the stress-energy term in Einstein's field equations would be zero everywhere, even when energy is present.

We will call this hypothetical infinitely rigid spacetime, characterized by the constant volume vacuum state $\mathring{\mathcal{V}} = \mathring{\mathcal{V}}_0$, the **null-gravity spacetime**.

5.3 Vacuum State $\mathring{\mathcal{V}}_+$

Definition: $\mathring{\mathcal{V}}_+ \equiv \{q \in \mathring{\mathcal{V}} \mid \Omega_{\max} > \Omega_t(q)\}$.

In this vacuum region we have a positive capacity to receive volumetric influx ($\Phi > 0$) everywhere, and no volumetric outflux. This describes a spacetime with innate slack. Volume will deform in response to energetic pressure, at least so long as the inserted energy's total volumetric displacement does not exceed the capacity of $\mathring{\mathcal{V}}_+$.

5.3.1 When $\mathring{\mathcal{V}} \rightarrow \mathring{\mathcal{V}}_+$

A vacuum spacetime where $\mathring{\mathcal{V}} = \mathring{\mathcal{V}}_+$ is innately capable of producing gravitational effects, and is thus the anticipated vacuum limit of our spacetime (the bottom line in fig. 8).




Figure 8: Vacuum States

5.4 Our Vacuum Spacetime

Due to the inherent instability of $\mathring{\mathcal{V}}_-$ shown in eq. (28), we expect $\Omega_t \leq \Omega_{\max}$ to be generically true across our spacetime, and hence propose that the volume of $\mathcal{M}$'s vacuum limit satisfies

$$\mathring{\mathcal{V}} = \overline{\mathring{\mathcal{V}}_+ \cup \mathring{\mathcal{V}}_0}. \quad (29)$$

We do allow for the possibility that Ω_t can exceed $\Omega_{\max}$ locally, and hence for $\dot{\mathcal{V}}_- \neq \emptyset$, though since eq. (28) shows no point of spacetime can be an element of $\dot{\mathcal{V}}_-$ for longer than the Planck time, we propose that $\dot{\mathcal{V}}_-$ is nowhere dense, and conjecture a likely relationship between its instability and the quantum vacuum fluctuations.

5.5 The Homogeneous Vacuum

To understand the homogeneous vacuum, we imagine the limit where $\dot{\mathcal{V}}$'s volume form Ω_t approaches constant value $\dot{\Omega}$. But first we will establish whether it makes physical sense for Ω_t to have the same volume across all of $\mathcal{V}$.

5.5.1 Compactness of $\mathcal{V}_+ \cup \mathcal{V}_-$

If we can correlate the influx eq. (25) to the observed cosmological distribution of dark energy, we will have direct observational evidence of volumetric flow. This will demonstrate that both the influx equation's denominator and its total gradient are finite and non-zero. A finite gradient demonstrates that the amount of energy causing volumetric distortions across the spacelike hypersurface is finite. A measurable gradient demonstrates that the portion of the hypersurface capable of receiving volumetric flux, $\mathcal{V}_+$, is also finite. Together these demonstrate that the portion of the universe which contains energy as well as that which is capable of responding to volumetric pressure from energy, are both finite and compact.

5.5.2 Homogeneous Vacuum Volume

If there exists a homogeneous vacuum across which every point has the same volume $\dot{\Omega}$ then we can simplify eq. (9) to

$$\mathbb{V} = \dot{\Omega} \int_{\mathcal{V}} \omega_0 \quad (30)$$

and it would then follow that $\mathbb{V}$ is finite only if $\int_{\mathcal{V}} \omega_0$ is finite and therefore only if $\mathcal{V}$ is compact. With the compactness of both $\mathcal{V}_+$ and $\mathcal{V}_-$ established, the only open question is whether $\mathcal{V}_0$ is compact. This means that the only open question is whether the portion of the universe where there is no energy and no ability to be impacted by energy's volumetric pressure—and hence where nothing energetic can happen aside from quantum background fluctuations—is compact. We defer to Archimedes' Sandreckoner a second time and propose $\mathcal{V}$ to be compact. We reason that with the finiteness of the totality of the energetic universe established, the alternative proposition—that the portion of the universe devoid of energetic interactions is infinite—is an extraordinary claim that lacks sufficient supporting evidence.

With compactness, we can define the homogeneous vacuum to be not only the limit where $\rho \rightarrow 0$ but also where Ω_t approaches the **homogeneous vacuum volume** $\dot{\Omega}$ at all points of $\mathcal{V}$. In addition, since $\dot{\Omega}$ must obey the same relations as

Ω_t , we can say that the set of physically meaningful choices for $\mathring{\Omega}$ is constrained by

$$0 < \mathring{\Omega} < \Omega_{\max} . \quad (31)$$

Finally, since $\mathring{\Omega} = \Omega_{\max}$ represents section 5.2's null-gravity spacetime, we can conclude that across the class of physically meaningful spacetimes for $\mathcal{M}$, as $\mathring{\Omega} \rightarrow \Omega_{\max}$, the homogeneous vacuum $\mathring{\mathcal{V}}$ will approach the null-gravity spacetime where $\mathring{\mathcal{V}}_+$ and $\mathring{\mathcal{V}}_-$ are both empty sets and $\mathring{\mathcal{V}} = \mathring{\mathcal{V}}_0$.

6 Gravitation

A full account of how the field equations of general relativity relate to this framework will be part of future work. For now, we show how the gravitational coupling relates to the framework and derive some immediate implications.

6.1 Gravitational Coupling

The observation that $\mathcal{V}$ approaches the null-gravity spacetime as $\mathring{\Omega} \rightarrow \Omega_{\max}$ in section 5.5.2 yields the important conclusion that a given spacetime's ability to deform in response to energy, and hence, the base strength of the gravitational coupling $\mathring{G}$, is proportional to the innate volumetric slack, $\Omega_{\max} - \mathring{\Omega}$, of its homogeneous vacuum volume, $\mathring{\mathcal{V}}$, and thus

$$\mathring{G} \propto \int_{\mathring{\mathcal{V}}} (\Omega_{\max} - \mathring{\Omega}) \omega_0 . \quad (32)$$

For a non-vacuum spacetime, we can replace $\Omega_{\max}$ with Ω_ρ , and $\mathring{\Omega}$ with Ω_t and conjecture

$$G_t \propto \int_{\mathcal{V}} (\Omega_\rho - \Omega_t) \omega_0 \quad (33)$$

for energetic spacetime's gravitational coupling, which is also the integral of the volumetric potential from eq. (15) across the universe's entire volume,

$$G_t \propto \int_{\mathcal{V}} \Phi \omega_0 . \quad (34)$$

To get a relation between G_t and $\mathring{G}$, we can insert $\Omega_{\max} - \Omega_{\max} = 0$ into eq. (33) and get

$$G_t \propto \int_{\mathcal{V}} (\Omega_{\max} - \Omega_t) \omega_0 - \int_{\mathcal{V}} (\Omega_{\max} - \Omega_\rho) \omega_0 , \quad (35)$$

then linearly adjust the scale of Ω_t to an equivalent form $\mathring{\Omega}_t$ which expresses the current volume as a perturbation from the homogenous vacuum volume $\mathring{\Omega}$ through

$$\mathring{\Omega}_t \equiv \Omega_t - \mathring{\Omega}. \quad (36)$$

Substituting this into eq. (35), we get

$$G_t \propto \int_{\mathcal{V}} (\Omega_{\max} - \mathring{\Omega}) \omega_0 - \int_{\mathcal{V}} \mathring{\Omega}_t \omega_0 - \int_{\mathcal{V}} (\Omega_{\max} - \Omega_\rho) \omega_0. \quad (37)$$

Finally, since

$$\int_{\mathcal{V}} \mathring{\Omega}_t \omega_0 = \int_{\mathcal{V}} (\Omega_t - \mathring{\Omega}) \omega_0 = \int_{\mathcal{V}} \Omega_t \omega_0 - \int_{\mathcal{V}} \mathring{\Omega} \omega_0 = \mathbb{V} - \mathbb{V} = 0, \quad (38)$$

we can zero-out the integral of $\mathring{\Omega}_t$ and substitute in the homogeneous vacuum's coupling from eq. (32).

$$G_t \propto \mathring{G} - \int_{\mathcal{V}} (\Omega_{\max} - \Omega_\rho) \omega_0. \quad (39)$$

6.2 Energetic Capacity

From eq. (33), we can see that a second null-gravity bound for $\mathcal{V}$ occurs if the volumetric limit from eq. (12) equals the available volume, and thus $\Omega_\rho = \Omega_t$ everywhere. As with the null-gravity spacetime (where $\mathring{\mathcal{V}}_+$ is empty), if no point in energetic spacetime has excess volumetric potential, then $\mathcal{V}_+$ will be empty, and $\mathcal{V}$ will have no capacity for accepting volumetric influx.

For a volume that has reached its energetic capacity, we can substitute eq. (12) into $\Omega_\rho = \Omega_t$ and get

$$\zeta \circ \rho_{\text{cap}} = \Omega_t. \quad (40)$$

Our proposition that the energy-volume relation ζ be smooth and strictly monotonic ensures the existence of an inverse. We can thus write

$$\rho_{\text{cap}} = \zeta^{-1} \circ \Omega_t \quad (41)$$

and integrate across the total volume to produce the **energetic capacity** of a volume form Ω_t , given by

$$\int_{\mathcal{V}} \rho_{\text{cap}} \omega = \int_{\mathcal{V}} (\zeta^{-1} \circ \Omega_t) \omega = \int_{\mathcal{V}} \Omega_t \cdot (\zeta^{-1} \circ \Omega_t) \omega_0. \quad (42)$$

If the total energy of the universe $\mathbb{E}$ becomes equal to or greater than the energetic capacity, then space becomes infinitely rigid as $\mathcal{V}_+ \rightarrow \emptyset$, and $G_t \rightarrow 0$. We conclude that for spacetime to exhibit gravitational effects, its total energy must remain bounded by

$$\mathbb{E} < \int_{\mathcal{V}} \Omega_t \cdot (\zeta^{-1} \circ \Omega_t) \omega_0. \quad (43)$$

For the homogeneous vacuum, this reduces to

$$\mathbb{E} < \zeta^{-1}(\mathring{\Omega}) \cdot \mathring{\Omega} \int_{\mathcal{V}} \omega_0, \quad (44)$$

We can then substitute in from eq. (30), and get an expression for the energetic capacity of the homogeneous vacuum that depends only on the total volume $\mathbb{V}$, the homogeneous vacuum volume $\mathring{\Omega}$, and the energy-volume relation ζ .

$$\frac{\mathbb{E}}{\mathbb{V}} < \zeta^{-1}(\mathring{\Omega}). \quad (45)$$

6.3 A Brief Investigation of ζ

We have specified throughout this paper that both the energy-volume relation ζ and the volume form Ω_t are positive definite and smooth. Though we can make a stronger statement regarding positive definiteness. Since the smallest resolvable non-zero volume is ℓ_{P}^3 , requiring $\Omega_t > 0$ is equivalent to requiring $\Omega_t \geq \ell_{\text{P}}^3$. More importantly, to prevent volumetric flow from driving Ω_t to a non-positive value, Φ must be non-negative whenever Ω_t equals its minimum $\Omega_t = \ell_{\text{P}}^3$, from which it follows that ζ also has a minimum value of ℓ_{P}^3 .

Though we expect ζ 's full form to have non-linear terms, we start by exploring a linear ζ to determine the slope of its linear approximation. The strictly monotonically decreasing property ensures the existence of ρ_{max} such that $\zeta(\rho_{\text{max}}) = \ell_{\text{P}}^3$. Solving for this condition yields the linear approximation,

$$\zeta(\rho)/\Omega_{\text{max}} = 1 - \frac{1 - \ell_{\text{P}}^3/\Omega_{\text{max}}}{\rho_{\text{max}}} \rho + \mathcal{O}(\rho^2). \quad (46)$$

Though, since the $\ell_{\text{P}}^3/\Omega_{\text{max}}$ term in the numerator is multiple orders of magnitude smaller than the Planck volume and thus unmeasurable and effectively zero, we can safely ignore the term and write

$$\zeta(\rho)/\Omega_{\text{max}} = 1 - \frac{1}{\rho_{\text{max}}} \rho + \mathcal{O}(\rho^2). \quad (47)$$

We use this form for ζ to highlight the obvious parallels with eq. (10) from general relativity. While our linear coefficient is $1/\rho_{\max}$ rather than the $(8\pi/15) \cdot (r^2 G/c^2)$ of general relativity, it is important to recognize that we cannot directly equate the two. Equation (10) is a series expansion in the radius around a point, while eq. (47) is a series expansion in the energy density around the vacuum.

6.3.1 A Closed Form

Conjecture: The energy volume relation ζ is infinitely flat as $\rho \rightarrow \infty$, with form

$$\zeta(\rho) = (\Omega_{\max} - \ell_{\text{P}}^3) \cdot e^{-\rho/\rho_{\max}} + \ell_{\text{P}}^3, \quad (48)$$

which is equivalent to

$$\zeta(\rho) = \Omega_{\max} \cdot e^{-\rho/\rho_{\max}} \quad (49)$$

except in regions of space where $\rho \gg \rho_{\max}$.

With this form for ζ , we can derive its inverse

$$\zeta^{-1}(\Omega) = \rho_{\max} \cdot \ln \left(\frac{\Omega_{\max}}{\Omega} \right) \quad (50)$$

and get a final form for the energetic capacity of the homogeneous vacuum from eq. (45)

$$\frac{\mathbb{E}}{\mathbb{V}} < \rho_{\max} \cdot \ln \left(\frac{\Omega_{\max}}{\Omega} \right). \quad (51)$$

Similarly, we get updated expressions for the volumetric potential from eq. (15),

$$\Phi = \Omega_{\max} \cdot e^{-\rho/\rho_{\max}} - \Omega. \quad (52)$$

and the gravitational coupling from eq. (34),

$$G_t \propto \int_{\mathcal{V}} \left(\Omega_{\max} \cdot e^{-\rho/\rho_{\max}} - \Omega \right) \omega_0. \quad (53)$$

Singularities

While numerically unimportant, the last ℓ_{P}^3 term in eq. (48) has theoretical importance. It ensures that the energy volume relation asymptotically approaches the Planck volume rather than 0. This protects Ω_t 's positive definiteness, and excludes the possibility of gravitational singularities occurring on the manifold.

The conjecture also aligns precisely with our preceding derivation, as eq. (47) is the first order Taylor series approximation of eq. (49).

7 Future Work

Numerical and Experimental Verification

There is a wealth of astronomical data available against which we can test our framework’s volume conservation mechanics. Relevant to dark matter, the spatial offset observed between the Bullet Cluster’s baryonic center of mass and its gravitational well[26, 27] could be a natural consequence of volumetric flow in the wake of a violent collision that significantly shifts a cluster’s center of mass away from the center of its residual stress. Similarly, numerical simulations of $\partial_t\Omega$ from section 4.3 could be tested against the Milky Way rotation data from the European Space Agency’s Gaia satellite[28–30]. Relevant to dark energy, the Dark Energy Spectroscopic Instrument’s mapping of baryon acoustic oscillations[31, 32] could have signatures consistent with eq. (25).

General Relativity

An examination of Einstein’s papers from the period between special and general relativity reveals early attempts to frame gravity as a scalar field conceptually similar to our framework[33]. The scalar field was set aside because the full richness of relativistic mechanics requires the metric tensor. However, our framework suggests that the idea of an underlying scalar field has merit and can coexist with the metric tensor in a full theory of gravity, by placing the scalar volume form at the foundation and presuming that the unimodular metric rests atop it.

With respect to existing relativity theory, the adjustment we make is to replace spacetime’s perfect elasticity with a locally plastic medium which continuously reflatens over time through a global volume conservation process analogous to the mechanical annealing of amorphous metallic glasses[34–36]. Aside from those cases where future investigation reveals that current theory’s topological assumptions are inconsistent with the volume conservation mechanics, we propose that relativistic dynamics in this framework behave as current theory dictates. For example, we expect that gravitational waves will propagate with the same volume-preserving shear that current theory predicts, while future work on the residual stress implications may reveal second order divergences from existing theory in the wave dynamics.

Significant theoretical work remains to complete a bridge to the tensor calculus of Einstein’s field equations. A specific future focus will address eq. (49)’s impact on cosmic inflation and black hole dynamics, where the higher order terms in ζ ’s Taylor approximation will dominate.

Quantum Field Theory

We believe that conjoining $\mathcal{Q}$ with 3-space by considering all points on the same spacelike hypersurface as the singular neighborhood of an indiscrete topological substrate, can serve as a useful model for framing the fields of quantum theory. One proposition is that the volume form Ω_t from this paper is a scalar quantum field, that this field provides coupled particles with the scale that $\tilde{\gamma}_{ij}$ lacks, and that the rate at which Ω_t samples Ω_ρ 's energetic pressure provides the tick of the Planck time. The sampling interval t_P and the local reference for the Planck scale ℓ_P then combine to determine light's speed $c = \ell_P/t_P$.

8 Acknowledgements

I am especially thankful to Henk Neefs for multiple inspiring discussions over breakfast in the Intel RNB Cafe. Additional thanks to Janna Levin's *How the Universe got its Spots*[37] for its characterization of late 1990s UC Berkeley physics culture. Reading it for the first time in 2026 stirred memories of my own time as an undergraduate in the physics department earlier the same decade. It was then that I first formed the thought that there must be something about the topological structure of space we are missing. A return to that time with the perspective of hindsight, and with my own poetic ruminations on the topic fully formed[38], is where the current paper started.

References

1. Einstein A (1965) On a heuristic point of view concerning the production and transformation of light. *American Journal of Physics* 33:367–374. <https://doi.org/10.1119/1.1971542>
2. Einstein A (1989) On the development of our views concerning the nature and constitution of radiation. In: *The collected papers of albert einstein*, vol. 2. Princeton University Press, pp 379–394
3. Broglie L de (1925) On the theory of quanta. *Annales de Physique* 10:22–128
4. Davisson C, Germer LH (1927) Diffraction of electrons by a crystal of nickel. *Phys Rev* 30:705–740. <https://doi.org/10.1103/PhysRev.30.705>
5. Shibata M, Nakamura T (1995) Evolution of three-dimensional gravitational waves: Harmonic slicing case. *Phys Rev D* 52:5428–5444
6. Baumgarte TW, Shapiro SL (1998) On the numerical integration of einstein's field equations. *Phys Rev D* 59:024007

7. Aspect A, Dalibard J, Roger G (1982) Experimental test of bell's inequalities using time-varying analyzers. *Phys Rev Lett* 49:1804–1807
8. Weihs G, Jennewein T, Simon C, et al (1998) Violation of bell's inequality under strict einstein locality conditions. *Phys Rev Lett* 81:5039–5043
9. Hensen B et al (2015) Loophole-free bell inequality violation using electron spins separated by 1.3 kilometres. *Nature* 526:682–686
10. Giustina M et al (2015) Significant-loophole-free test of bell's theorem with entangled photons. *Phys Rev Lett* 115:250401
11. Shalm LK et al (2015) Strong loophole-free test of local realism. *Phys Rev Lett* 115:250402
12. Steinberg AM, Kwiat PG, Chiao RY (1993) Measurement of the single-photon tunneling time. *Phys Rev Lett* 71:708–711
13. Sainadh US, Xu H, Wang X, et al (2017) [Attosecond angular streaking and tunnelling time in atomic hydrogen](#)
14. Ramos R, Spierings D, Racicot I, Steinberg AM (2020) Measurement of the time spent by a tunnelling atom within the barrier region. *Nature* 583:529–532
15. Einstein A (2009) Quantum theory at the crossroads: Reconsidering the 1927 solvay conference. In: Bacciagaluppi G, Valentini A (eds). Cambridge Univ. Press, pp 440–442
16. Einstein A, Podolsky B, Rosen N (1935) Can quantum-mechanical description of physical reality be considered complete? *Phys Rev* 47:777–780
17. Bell JS (1964) On the einstein podolsky rosen paradox. *Physics Physique Fizika* 1:195–200
18. Maldacena J, Susskind L (2013) Cool horizons for entangled black holes. *Fortsch Phys* 61:781–811
19. Gray A, Vanhecke L (1979) Riemannian geometry as determined by the volumes of small geodesic balls. *Acta Math* 142:157–198
20. Feoli A (2018) Plasticity of spacetime as an alternative to dark matter. *Mod Phys Lett A* 33:1850194

21. Archimedes (1956) The sand reckoner. In: Newman JR (ed) The world of mathematics vol. 1. Simon; Schuster, pp 420–429
22. Dyson FW, Eddington AS, Davidson C (1920) A determination of the deflection of light by the sun’s gravitational field, from observations made at the total eclipse of may 29, 1919. Philosophical Transactions of the Royal Society of London Series A 220:291–333
23. Bertotti B, Iess L, Tortora P (2003) A test of general relativity using radio links with the cassini spacecraft. Nature 425:374–376
24. Rudin W (1976) Principles of mathematical analysis, 3rd ed. McGraw-Hill
25. Planck M (2024) [On the law of the energy distribution in the normal spectrum](#)
26. Clowe D, Bradač M, Gonzalez AH, et al (2006) A direct empirical proof of the existence of dark matter. The Astrophysical Journal Letters 648:L109–L113. <https://doi.org/10.1086/508162>
27. Markevitch M, Gonzalez AH, Clowe D, et al (2004) Direct constraints on the dark matter self-interaction cross section from the merging galaxy cluster 1E 0657-56. The Astrophysical Journal 606:819–824. <https://doi.org/10.1086/383158>
28. Jiao Y, Hammer F, Wang H, et al (2023) Detection of the Keplerian decline in the Milky Way rotation curve. Astronomy & Astrophysics 678:A208. <https://doi.org/10.1051/0004-6361/202347513>
29. Moffat JW (2024) [Implications of the Milky Way Declining Rotation Curve](#). arXiv e-prints
30. Melchiorri A et al (2026) [The Rotation Curve of the Milky Way: State of the Art, the Keplerian decline debate, and implications for dark matter](#). arXiv e-prints
31. Adame AG et al (2024) [DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations](#). arXiv e-prints
32. DESI Collaboration (2025) [Dynamical Dark Energy in light of the DESI DR2](#). arXiv e-prints

- 33. Einstein A (1996) The speed of light and the statics of the gravitational field. In: The collected papers of albert einstein, volume 4: The swiss years: Writings, 1912–1914 (english translation supplement). Princeton University Press, Princeton, NJ, pp 95–106
- 34. Dreyer O (2012) Internal relativity. *Foundations of Physics* 42:1346–1361. <https://doi.org/10.1007/s10701-012-9658-0>
- 35. Dreyer O (2008) [Early universe cosmology in internal relativity](#). arXiv preprint
- 36. Spaepen F (1977) A microscopic mechanism for steady state inhomogeneous flow in metallic glasses. *Acta Metallurgica* 25:407–415. [https://doi.org/10.1016/0001-6160\(77\)90232-2](https://doi.org/10.1016/0001-6160(77)90232-2)
- 37. Levin J (2002) How the universe got its spots: Diary of an unobservable universe. Princeton University Press, Princeton, NJ
- 38. Chamberlain JD (2025) [Symmetry emergent](#)